

GSK3β (Phospho Ser9) Rabbit pAb

CatalogNo: YP0124 Orthogonal Validated O

Key Features

Host Species • Rabbit	Reactivity Human,Mouse,Rat,Drosophila 	Applications IF,WB,IHC,IP,ELISA
MW • 48kD (Observed)	lsotype • lgG	

Recommended Dilution Ratios

IF 1:50-200 WB 1:500-1:2000 IHC 1:100-1:300 IP 2-5 ug/mg lysate ELISA 1:5000 Not yet tested in other applications

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human GSK3
beta around the phosphorylation site of Ser9. AA range:1-50

Specificity

Phospho-GSK3 β (S9) Polyclonal Antibody detects endogenous levels of GSK3 β protein only when phosphorylated at S9.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TTsFA

Target Information

GSK3B Gene name **Protein Name** Glycogen synthase kinase-3 beta Organism **Gene ID UniProt ID** Human 2932; P49841; Mouse Q9WV60; 56637; Rat 84027: P18266;

- CellularCytoplasm . Nucleus . Cell membrane . The phosphorylated form shows localization to
cytoplasm and cell membrane (PubMed:20937854). The MEMO1-RHOA-DIAPH1 signaling
pathway controls localization of the phosphorylated form to the cell membrane
(PubMed:20937854). .
- **Tissue specificity** Expressed in testis, thymus, prostate and ovary and weakly expressed in lung, brain and kidney. Colocalizes with EIF2AK2/PKR and TAU in the Alzheimer disease (AD) brain.
- Function Catalytic activity:ATP + [tau protein] = ADP + [tau protein] phosphate.,enzyme regulation:Inhibited when phosphorylated by AKT1.,Function:Participates in the Wnt signaling pathway. Implicated in the hormonal control of several regulatory proteins including glycogen synthase, MYB and the transcription factor JUN. Phosphorylates JUN at sites proximal to its DNA-binding domain, thereby reducing its affinity for DNA. Phosphorylates MUC1 in breast cancer cells, and decreases the interaction of MUC1 with CTNNB1/beta-catenin.,PTM:Phosphorylated by AKT1 and ILK1.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. GSK-3 subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Monomer (By similarity). Interacts with CABYR, MUC1, NIN and PRUNE.,tissue specificity:Expressed in testis, thymus, prostate and ovary and weakly expressed in lung, brain and kidney.,

Validation Data



Perturbases SW1990
 Perturbases SW199



Ge, Lu, et al. "PRMT5 promotes epithelial-mesenchymal transition via EGFR- β -catenin axis in pancreatic cancer cells." Journal of cellular and molecular medicine 24.2 (2020): 1969-1979.



Ou, Liping, et al. "Dickkopf Wnt signaling pathway inhibitor 1 regulates the differentiation of mouse embryonic stem cells in vitro and in vivo." Molecular medicine reports 13.1 (2016): 720-730.



Immunofluorescence analysis of rat-spleen tissue. 1,GSK3β (phospho Ser9) Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of rat-spleen tissue. 1,GSK3β (phospho Ser9) Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Rat-testis tissue. 1,GSK3 β (phospho Ser9) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only. Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1,GSK3β (phospho Ser9) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.

Immunohistochemical analysis of paraffin-embedded Mouse-liver tissue. 1,GSK3β (phospho Ser9) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.

tempeRature, 30min). Negative control was used by secondary antibody only. Immunohistochemical analysis of paraffin-embedded Mouse-colon tissue.

Immunohistochemical analysis of paraffin-embedded Mouse-colon tissue. 1,GSK3β (phospho Ser9) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.

Immunohistochemical analysis of paraffin-embedded Mouse-lung tissue. 1,GSK3 β (phospho Ser9) Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.

Western Blot analysis of Hela cells using Phospho-GSK3 β (S9) Polyclonal Antibody diluted at 1:1000

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using GSK3 beta (Phospho-Ser9) Antibody. The picture on the right is blocked with the phospho peptide.











Western blot analysis of lysates from HeLa cells treated with EGF, using GSK3 beta (Phospho-Ser9) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **GSK3β (Phospho Ser9) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents