eNOS (Phospho Thr494) Rabbit pAb

CatalogNo: YP0196

Key Features

Host Species	Reactivity	Applications
• Rabbit	• Human,Mouse,Rat	• WB,ELISA

MW	Isotype
• 130-140kD (Observed)	• IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000
ELISA 1:5000
Not yet tested in other applications.

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human eNOS around the phosphorylation site of Thr494. AA range:462-511
Specificity	Phospho-NOS3 (T494) Polyclonal Antibody detects endogenous levels of NOS3 protein only when phosphorylated at T494.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):KKtFK

Target Information

Gene name NOS3

Protein Name Nitric oxide synthase endothelial


Organism	Gene ID	UniProt ID
Human	4846;	P29474;
Mouse	18127;	P70313;
Rat	24600;	Q62600;

Cellular Localization Cell membrane. Membrane , caveola. Cytoplasm , cytoskeleton. Golgi apparatus. Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity.


Tissue specificity Platelets , placenta , liver and kidney.

Function Catalytic activity:L-arginine + n NADPH + n H (+) + m O (2) = citrulline + nitric oxide + n NADP (+) . ,cofactor:Binds 1 FAD. ,cofactor:Binds 1 FMN. ,cofactor:Heme group. ,cofactor:Tetrahydrobiopterin (BH4) . May stabilize the dimeric form of the enzyme. ,enzyme regulation:Stimulated by calcium/calmodulin. Inhibited by NOSIP and NOSTRIN. ,Function:Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF) -induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets. ,online information:Nitric oxide synthase entry ,polymorphism:Variation in NOS3 seem to be associated with susceptibility to coronary spasm. ,similarity:Belongs to the NOS family. ,similarity:Contains 1 FAD-binding FR-type domain. ,similarity:Contains 1 flavodoxin-like domain. ,subcellular location:Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity. ,subunit:Homodimer. Interacts with NOSIP and NOSTRIN. ,tissue specificity:Platelets , placenta , liver and kidney. ,


Validation Data



Western Blot analysis of 293 cells using Phospho-NOS3 (T494) Polyclonal Antibody diluted at 1:1000



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using eNOS (Phospho-Thr494) Antibody



Western blot analysis of lysates from Jurkat cells, using eNOS (Phospho-Thr494) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
eNOS (Phospho Thr494) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.