STAT3 (Phospho Tyr705) Rabbit pAb

CatalogNo: YP0251 Orthogonal Validated Comparable Abs

Key Features

Host Species
- Rabbit

Reactivity
- Human,Mouse,Rat,Pig(Testbyoutcustomer)

Applications
- WB,IHC,IF,IP,ELISA

MW
- 88kD (Observed)

Isotype
- IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IF 1:50-200
WB 1:500-1:2000
IHC 1:100-1:300
IP 2-5 ug/mg lysate
ELISA 1:20000
Not yet tested in other applications

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human STAT3 around the phosphorylation site of Tyr705. AA range:672-721

Specificity	Phospho-Stat3 (Y705) Polyclonal Antibody detects endogenous levels of Stat3 protein only when phosphorylated at Y705.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):APyLK

Target Information

Gene name	STAT3 APRF
Protein Name	Signal transducer and activator of transcription 3

Organism	Gene ID	UniProt ID
Human	6774;	P40763;
Mouse	20848;	P42227;
Rat	25125;	P52631;

Cellular Localization	Cytoplasm . Nucleus . Shuttles between the nucleus and the cytoplasm. Translocated into the nucleus upon tyrosine phosphorylation and dimerization , in response to signaling by activated FGFR1 , FGFR2 , FGFR3 or FGFR4. Constitutive nuclear presence is independent of tyrosine phosphorylation. Predominantly present in the cytoplasm without stimuli. Upon leukemia inhibitory factor (LIF) stimulation , accumulates in the nucleus. The complex composed of BART and ARL2 plays an important role in the nuclear translocation and retention of STAT3. Identified in a complex with LYN and PAG1.
Tissue specificity	Heart , brain , placenta , lung , liver , skeletal muscle , kidney and pancreas. Expressed in naive CD4 (+) T cells as well as T-helper Th17 , Th1 and Th2 cells (PubMed:31899195) .

Function	Disease:Defects in STAT3 are the cause of hyperimmunoglobulin E recurrent infection syndrome autosomal dominant (AD-HIES) [MIM:147060]; also known as hyper-IgE syndrome or Job syndrome. AD-HIES is a rare disorder of immunity and connective tissue characterized by immunodeficiency , chronic eczema , recurrent Staphylococcal infections , increased serum IgE , eosinophilia , distinctive coarse facial appearance , abnormal dentition , hyperextensibility of the joints , and bone fractures. ,Function:Transcription factor that binds to the interleukin-6 (IL-6) -responsive elements identified in the promoters of various acute-phase protein genes. Activated by IL31 through IL31RA. ,miscellaneous:Involved in the gp130-mediated signaling pathway. ,online information:STAT3 entry ,online information:STAT3 mutation db ,PTM:Tyrosine phosphorylated in response to IL-6 , IL-11 , CNTF , LIF , CSF-1 , EGF , PDGF , IFN-alpha and OSM. Phosphorylated on serine upon DNA damage , probably by ATM or ATR. Serine phosphorylation is important for the formation of stable DNA-binding STAT3 homodimers and maximal transcriptional activity. ,similarity:Belongs to the transcription factor STAT family. ,similarity:Contains 1 SH2 domain. ,subcellular location:Shuttles between the nucleus and the cytoplasm. Constitutive nuclear presence is independent of tyrosine phosphorylation. ,subunit:Forms a homodimer or a heterodimer with a related family member (at least STAT1) . Interacts with NCOA1 , PELP1 , SOCS7 and STATIP1. Interacts with HCV core protein. Interacts with IL23R in presence of IL23. Interacts with IL31RA. Interacts with SIPAR. Interacts (via SH2 domain) with NLK (By similarity) . Interacts with KPNA4 and KPNA5; KPNA4 may be the primary mediator of nuclear import (By similarity) . Interacts with TMF1. ,tissue specificity:Heart , brain , placenta , lung , liver , skeletal muscle , kidney and pancreas. ,

Validation Data

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
STAT3 (Phospho Tyr705) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.