

STAT5A/B (Phospho Tyr694/699) Rabbit pAb

CatalogNo: YP0254

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 91kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human STAT5A around the phosphorylation site of Tyr694. AA range: 666-715

Specificity This antibody detects endogenous levels of STAT5A/B only when phosphorylated at Human:Y694/699, Mouse:Y694/699, Rat:Y694/699..The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): DGyVK

Target Information

Gene name	STAT5A/STAT5B		
Protein Name	Signal transducer and activator of transcription 5A/B		
	Organism	Gene ID	UniProt ID
	Human	6776 ; 6777 ;	P42229 ; P51692 ;
	Mouse	20850 ; 20851 ;	
	Rat	24918 ; 25126 ;	Q62771 ; P52632 ;
Cellular Localization	Cytoplasm . Nucleus . Translocated into the nucleus in response to phosphorylation.		
Tissue specificity	Brain,Cervix,Epithelium,Lung,Placenta,Synovial memb		
Function	Function:Carries out a dual Function: signal transduction and activation of transcription. Binds to the GAS element and activates PRL-induced transcription.,online information:STAT5 entry,PTM:Tyrosine phosphorylated in response to IL-2, IL-3, IL-7, IL-15, GM-CSF, growth hormone, prolactin, erythropoietin and thrombopoietin. Tyrosine phosphorylation is required for DNA-binding activity and dimerization. Serine phosphorylation is also required for maximal transcriptional activity.,similarity:Belongs to the transcription factor STAT family.,similarity:Contains 1 SH2 domain.,subcellular location:Translocated into the nucleus in response to phosphorylation.,subunit:Forms a homodimer or a heterodimer with a related family member. Binds NR3C1 (By similarity). Interacts with NCOA1 and SOCS7.,		

Validation Data

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
STAT5A/B (Phospho Tyr694/699) Rabbit pAb