

GABAA Rβ1 (Phospho Ser434) Rabbit pAb

CatalogNo: YP0576

Key Features

Host Species • Rabbit	Reactivity Human,Mouse,Rat,Monkey 	Applications • WB,IHC,IF,ELISA
MW • 55kD (Observed)	Isotype • IgG	

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000 IF 1:50-200

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human GABA-RB
around the phosphorylation site of Ser434. AA range:401-450

Specificity Phospho-GABAA Rβ1 (S434) Polyclonal Antibody detects endogenous levels of GABAA Rβ1 protein only when phosphorylated at S434.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RAsQL

Target Information

Gene name	GABRB1
-----------	--------

Protein Name


Gamma-aminobutyric acid receptor subunit beta-1

Organism	Gene ID	UniProt ID
Human	<u>2560;</u>	<u>P18505;</u>
Mouse	<u>14400;</u>	<u>P50571;</u>
Rat	<u>25450;</u>	<u>P15431;</u>


Cellular Cell junction, synapse, postsynaptic cell membrane ; Multi-pass membrane protein . Cell Localization membrane ; Multi-pass membrane protein .

Tissue specificity Brain, Epithelium,


Function Function:GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel.,online information:Forbidden fruit - Issue 56 of March 2005, similarity: Belongs to the ligand-gated ionic channel (TC 1.A.9) family., subunit: Binds UBQLN1 (By similarity). Generally pentameric. There are five types of GABA(A) receptor chains: alpha, beta, gamma, delta, and rho.,



Western Blot analysis of various cells using Phospho-GABAA R^β1 (S434) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GABA-RB (Phospho-Ser434) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using GABA-RB (Phospho-Ser434) Antibody. The picture on the right is blocked with the phospho peptide.

	117 85
GABA-RB (pSer434)	48 34
	26
	19 (kD)

Western blot analysis of lysates from COS7 cells, using GABA-RB (Phospho-Ser434) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: GABAA Rβ1 (Phospho Ser434) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents