

Synaptotagmin 1/2 (Phospho Thr202/199) Rabbit pAb

CatalogNo: YP0587

Orthogonal Validated 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 60kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-1:2000**IHC 1:100-1:300****ELISA 1:40000****IF 1:50-200**

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information


Immunogen The antiserum was produced against synthesized peptide derived from human Synaptotagmin around the phosphorylation site of Thr202. AA range: 176-225

Specificity This antibody detects endogenous levels of Synaptotagmin 1/2 only when phosphorylated at Human:T202/199, Mouse:T201/202, Rat:T201/202..The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RKtLN


Target Information

Gene name	SYT1/SYT2		
Protein Name	Synaptotagmin-1/2		
	Organism	Gene ID	UniProt ID
	Human	6857 ; 127833 ;	P21579 ; Q8N9I0 ;
	Mouse	20979 ; 20980 ;	
	Rat	25716 ; 24805 ;	P21707 ; P29101 ;
Cellular Localization	Cytoplasmic vesicle, secretory vesicle membrane ; Single-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Single-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, chromaffin granule membrane ; Single-pass membrane protein . Cytoplasm .		
Tissue specificity	Expressed in melanocytes (PubMed:23999003).		
Function	cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domains.,Domain: The first C2 domain mediates Ca(2+)-dependent phospholipid binding.,Domain: The second C2 domain mediates interaction with SV2A and STN2.,Function: May have a regulatory role in the membrane interactions during trafficking of synaptic vesicles at the active zone of the synapse. It binds acidic phospholipids with a specificity that requires the presence of both an acidic head group and a diacyl backbone. A Ca(2+)-dependent interaction between synaptotagmin and putative receptors for activated protein kinase C has also been reported. It can bind to at least three additional proteins in a Ca(2+)-independent manner; these are neurexins, syntaxin and AP2.,similarity: Belongs to the synaptotagmin family.,similarity: Contains 2 C2 domains.,subcellular location: Synaptic vesicles and chromaffin granules.,subunit: Homotetramer (Probable). Interacts with SCAMP5, STN2, SV2A, SV2B, SV2C and RIMS1. Forms a complex with SV2B, syntaxin 1 and SNAP25.,		


Validation Data




Western Blot analysis of various cells using Phospho-Synaptotagmin 1/2 (T202/199) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Synaptotagmin (Phospho-Thr202) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Synaptotagmin (Phospho-Thr202) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with Forskolin 40nM 30', using Synaptotagmin (Phospho-Thr202) Antibody. The lane on the left is blocked with the phospho peptide.

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**Synaptotagmin 1/2
(Phospho
Thr202/199) Rabbit
pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)