

c-Fms (Phospho Tyr561) Rabbit pAb

Reactivity

Isotype

IgG

CatalogNo: YP0734

Key Features

Host Species

Rabbit
 Human, Mouse, Rat

Applications
• WB,IHC,IF,ELISA

MW

• 130-170kD (Observed)

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:10000 IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human CSFR

around the phosphorylation site of Tyr561. AA range:531-580

Specificity Phospho-c-Fms (Y561) Polyclonal Antibody detects endogenous levels of c-Fms protein

only when phosphorylated at Y561. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):NSyTF

| Target Information

Gene name

CSF1R

Protein Name

Macrophage colony-stimulating factor 1 receptor

Organism	Gene ID	UniProt ID
Human	<u>1436;</u>	<u>P07333;</u>
Mouse	<u>12978;</u>	<u>P09581;</u>
Rat		<u>Q00495</u> ;

Cellular Localization

Cell membrane; Single-pass type I membrane protein.

Tissue specificity Expressed in bone marrow and in differentiated blood mononuclear cells.

Function


Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., Function: Protein tyrosine-kinase transmembrane receptor for CSF1 and IL34., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family.

CSF-1/PDGF receptor subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 5 Ig-like C2-type (immunoglobulin-like)


domains...subunit:Interacts with INPPL1/SHIP2 and THOC5..tissue specificity:Expressed in

bone marrow and in differentiated blood mononuclear cells.,

I Validation Data



Western Blot analysis of HepG2 cells using Phospho-c-Fms (Y561) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using CSFR (Phospho-Tyr561) Antibody. The picture on the right is blocked with the phospho peptide.

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: c-Fms (Phospho Tyr561) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents