Applications

WB,IHC,IF,ELISA



ATRIP (Phospho Ser224) Rabbit pAb

CatalogNo: YP0813 Orthogonal Validated 💽

Key Features

Host Species Reactivity

Rabbit
 Human,Rat,Mouse,

MW Isotype
• 86kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:10000 IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human ATRIP

around the phosphorylation site of Ser224. AA range:190-239

Specificity

Phospho-ATRIP (S224) Polyclonal Antibody detects endogenous levels of ATRIP protein only when phosphorylated at S224. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):HVsPR

| Target Information

Gene name

ATRIP

Protein Name

ATR-interacting protein

Orga	nism	Gene ID	UniProt ID
Hun	nan	<u>84126;</u>	Q8WXE1;
Mou	ıse		Q8BMG1;

Cellular Localization


Nucleus . Redistributes to discrete nuclear foci upon DNA damage.

Tissue specificity Ubiquitous.


Function

Caution:The gene for this protein is either identical to or adjacent to that of TREX1. Some of the mRNAs that encode ATRIP also encode TREX1 in another reading frame.,Domain:The EEXXXDDL motif is required for the interaction with catalytic subunit PRKDC and its recruitment to sites of DNA damage.,Function:Required for checkpoint signaling after DNA damage. Required for ATR expression, possibly by stabilizing the protein.,PTM:Phosphorylated by ATR.,sequence Caution:Translation N-terminally extended.,similarity:Belongs to the ATRIP family.,subcellular location:Redistributes to discrete nuclear foci upon DNA damage.,subunit:Heterodimer with ATR. The heterodimer binds the RPA complex and is then recruited to single stranded DNA. Interacts with CEP164 (via N-terminus).,tissue specificity:Ubiquitous.,

Validation Data



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using ATRIP (Phospho-Ser224) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with UV 15', using ATRIP (Phospho-Ser224) Antibody. The lane on the right is blocked with the phospho peptide.

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **ATRIP (Phospho**

ATRIP (Phospho Ser224) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents