

#### www.immunoway.com

## MAPKAPK-2 (Phospho Ser272) Rabbit pAb

CatalogNo: YP0819 Orthogonal Validated 💽

#### Key Features

Host Species	Reactivity	Applications
• Rabbit	• Human,Mouse,Rat	• WB,IHC,IF,ELISA
MW • 45-52kD (Observed)	Isotype • IgG	

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:5000 IF 1:50-200

#### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

ImmunogenThe antiserum was produced against synthesized peptide derived from human<br/>MAPKAPK2 around the phosphorylation site of Ser272. AA range:238-287

Specificity

Phospho-MAPKAPK-2 (S272) Polyclonal Antibody detects endogenous levels of MAPKAPK-2 protein only when phosphorylated at S272.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):AlsPG

### Target Information

Gene name MAPKAPK2

**Protein Name** MAP kinase-activated protein kinase 2

Organism	Gene ID	UniProt ID
Human	<u>9261;</u>	<u>P49137;</u>
Mouse	<u>17164;</u>	<u>P49138;</u>

# CellularCytoplasm . Nucleus . Phosphorylation and subsequent activation releases the<br/>autoinhibitory helix, resulting in the export from the nucleus into the cytoplasm.

**Tissue specificity** Expressed in all tissues examined.

**Function** Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Seems to be activated by two distinct pathways: the first involves the stimulation of p42/p44 MAPK by growth factors, the second, triggered by stress and heat shock, depends on the activation of MPK2 and upstream MAPKK/MAPKKK.,Function:Its physiological substrate seems to be the small heat shock protein (HSP27/HSP25). In vitro can phosphorylate glycogen synthase at 'Ser-7' and tyrosine hydroxylase (on 'Ser-19' and 'Ser-40'). This kinase phosphorylates Ser in the peptide sequence, Hyd-X-R-X(2)-S, where Hyd is a large hydrophobic residue (By similarity). Mediates both ERK and p38 MAPK/MAPK14 dependent neutrophil responses. Participates in TNF alpha-stimulated exocytosis of secretory vesicles in neutrophils. Plays a role in phagocytosis-induced respiratory burst activity.,PTM:Phosphorylated and activated by MAP kinase.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with PHC2.,tissue specificity:Expressed in all tissues examined.,

#### Validation Data



Immunohistochemistry analysis of paraffin-embedded human brain, using MAPKAPK2 (Phospho-Ser272) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from RAW264.7 cells treated with UV 15', using MAPKAPK2 (Phospho-Ser272) Antibody. The lane on the right is blocked with the phospho peptide.

### **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: MAPKAPK-2 (Phospho Ser272) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents