CXCR4 (Phospho Ser339) Rabbit pAb

CatalogNo: YP0917 Orthogonal Validated

Key Features

Host Species
- Rabbit

Reactivity
- Human,Mouse,Rat,Monkey

Applications
- WB,IHC,IF,ELISA

MW
- 38kD (Observed)

Isotype
- IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000
IHC 1:100-1:300
IF 1:200-1:1000
ELISA 1:20000
Not yet tested in other applications.

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human CXCR4 around the phosphorylation site of Ser339. AA range:303-352

Specificity	Phospho-Fusin (S339) Polyclonal Antibody detects endogenous levels of Fusin protein only when phosphorylated at S339.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):HSsVS

Target Information


Gene name	CXCR4
Protein Name	C-X-C chemokine receptor type 4

Organism	Gene ID	UniProt ID
Human	7852;	P61073;
Mouse	12767;	P70658;
Rat		O08565;


Cellular Localization	Cell membrane ; Multi-pass membrane protein . Cell junction. Early endosome. Late endosome. Lysosome. In unstimulated cells , diffuse pattern on plasma membrane. On agonist stimulation , colocalizes with ITCH at the plasma membrane where it becomes ubiquitinated. In the presence of antigen , distributes to the immunological synapse forming at the T-cell-APC contact area , where it localizes at the peripheral and distal supramolecular activation cluster (SMAC) .
Tissue specificity	Expressed in numerous tissues , such as peripheral blood leukocytes , spleen , thymus , spinal cord , heart , placenta , lung , liver , skeletal muscle , kidney , pancreas , cerebellum , cerebral cortex and medulla (in microglia as well as in astrocytes) , brain microvascular , coronary artery and umbilical cord endothelial cells. Isoform 1 is predominant in all tissues tested.

Function Alternative products:Additional isoforms seem to exist ,Caution:Was originally (PubMed:8329116 and PubMed:8234909) thought to be a receptor for neuropeptide Y type 3 (NPY3R) (NPY3-R) . ,Disease:Defects in CXCR4 are a cause of WHIM syndrome [MIM:193670]; also called warts , hypogammaglobulinemia , infections , and myelokathexis. WHIM syndrome is an immunodeficiency disease characterized by neutropenia , hypogammaglobulinemia and extensive human papillomavirus (HPV) infection. Despite the peripheral neutropenia , bone marrow aspirates from affected individuals contain abundant mature myeloid cells , a condition termed myelokathexis. ,Domain:The amino-terminus is critical for ligand binding. Residues in all four extracellular regions contribute to HIV-1 coreceptor activity. ,Function:Receptor for the C-X-C chemokine CXCL12/SDF-1. Transduces a signal by increasing the intracellular calcium ions level. Involved in haematopoiesis and in cardiac ventricular septum formation. Plays also an essential role in vascularization of the gastrointestinal tract , probably by regulating vascular branching and/or remodeling processes in endothelial cells. Could be involved in cerebellar development. In the CNS , could mediate hippocampal-neuron survival. Acts as a coreceptor (CD4 being the primary receptor) for HIV-1 X4 isolates and as a primary receptor for some HIV-2 isolates. Promotes Env-mediated fusion of the virus. ,online information:CXC chemokine receptors entry ,online information:CXCR4 entry ,online information:CXCR4 mutation db ,PTM:O- and N-glycosylated. Asn-11 is the principal site of N-glycosylation. There appears to be very little or no glycosylation on Asn-176. N-glycosylation masks coreceptor function in both X4 and R5 laboratory-adapted and primary HIV-1 strains through inhibiting interaction with their Env glycoproteins. The O-glycosylation chondroitin sulfate attachment does not affect interaction with CXCL12/SDF-1alpha nor its coreceptor activity. ,PTM:Sulfation on Tyr-21 is required for efficient binding of CXCL12/SDF-1alpha and promotes its dimerization. ,similarity:Belongs to the G-protein coupled receptor 1 family. ,subunit:Monomer. Can form dimers. Interacts with HIV-1 surface protein gp120 and Tat. ,tissue specificity:Expressed in numerous tissues , such as peripheral blood leukocytes , spleen , thymus , spinal cord , heart , placenta , lung , liver , skeletal muscle , kidney , pancreas , cerebellum , cerebral cortex and medulla (in microglia as well as in astrocytes) , brain microvascular , coronary artery and umbilical cord endothelial cells. Isoform 1 is predominant in all tissues tested. ,


Validation Data




Immunofluorescence analysis of Hela cell. 1,Fusin (phospho Ser339) Polyclonal Antibody (green) was diluted at 1:200 (4℃ overnight). 2, Goat Anti Rabbit AbFluor 488 Catalog:RS3211 was diluted at 1:1000 (room temperature, 50min). 3 DAPI (blue) 10min.




Western Blot analysis of HuvEc etoposide 25uM 24H cells using Phospho-Fusin (S339) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using CXCR4 (Phospho-Ser339) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using CXCR4 (Phospho-Ser339) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HUVEC cells treated with etoposide 25uM 24H, using CXCR4 (Phospho-Ser339) Antibody. The lane on the right is blocked with the phospho peptide.

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
CXCR4 (Phospho Ser339) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.