IFN-α/βRα (Phospho Tyr466) Rabbit pAb

CatalogNo: YP0989

Key Features

Host Species	Reactivity	Applications
• Rabbit	• Human,Mouse,Rat	• WB,IHC,IF,ELISA

MW	Isotype
• 63kD (Observed)	• IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000
IHC 1:100-1:300
ELISA 1:10000
IF 1:50-200

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human Interferon-alpha/beta Receptor alpha around the phosphorylation site of Tyr466. AA range:436-485

Specificity	Phospho-IFN-α/βRα (Y466) Polyclonal Antibody detects endogenous levels of IFN-α/βRα protein only when phosphorylated at Y466.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):INyVF


Target Information

Gene name	IFNAR1
Protein Name	Interferon alpha/beta receptor 1


Organism	Gene ID	UniProt ID
Human	3454;	P17181;
Mouse		P33896;

Cellular Localization	[Isoform 1]: Cell membrane ; Single-pass type I membrane protein . Late endosome . Lysosome . Interferon binding triggers internalization of the receptor from the cell membrane into endosomes and then into lysosomes. .
Tissue specificity	IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1 , isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266B1. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN-alpha resistant myeloma cell line U266R.
Function	Function:Receptor for interferons alpha and beta. Binding to type I IFNs triggers tyrosine phosphorylation of a number of proteins including JAKs , TYK2 , STAT proteins and IFNR alpha- and beta-subunits themselves. ,PTM:Phosphorylated on tyrosine residues by TYK2 tyrosine kinase. ,sequence Caution:Contaminating sequence. Potential poly-A sequence. ,similarity:Belongs to the type II cytokine receptor family. ,similarity:Contains 3 fibronectin type-III domains. ,tissue specificity:IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1 , isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266S. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R , isoform 1 is not expressed in U266R. ,

Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Interferon-alpha/beta Receptor alpha (Phospho-Tyr466) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using Interferon-alpha/beta Receptor alpha (Phospho-Tyr466) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
IFN-α/βRα (Phospho Tyr466) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.