

# Flk-1/Flt-4 (Phospho Tyr1054/1063) Rabbit pAb

CatalogNo: YP0996

### **Key Features**

**Host Species** Reactivity

 Rabbit Human, Mouse, Rat

Isotype

IgG

**Applications** IHC,IF,ELISA

MW 152kD (Calculated)

### Recommended Dilution Ratios

IHC 1:100-1:300 **ELISA 1:40000** IF 1:50-200

### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

The antiserum was produced against synthesized peptide derived from human VEGFR2 **Immunogen** 

around the phosphorylation site of Tyr1054. AA range:1020-1069

**Specificity** This antibody detects endogenous levels of VEGFR2/VEGFR3 only when phosphorylated

at Human:Y1054/Y1063, Mouse:Y1052/Y1063, Rat:Y1050/1063..The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):DIvKD

### **Target Information**

#### Gene name

KDR/FLT4

#### **Protein Name**

Vascular endothelial growth factor receptor 2/3

Organism	Gene ID	UniProt ID
Human	<u>3791; 2324;</u>	<u>P35968; P35916;</u>
Mouse	<u>14257;</u>	
Rat	25589; 114110;	<u>008775; Q91ZT1;</u>

### Cellular Localization


Cell junction. Endoplasmic reticulum. Cell membrane. Localized with RAP1A at cell-cell junctions (By similarity). Colocalizes with ERN1 and XBP1 in the endoplasmic reticulum in endothelial cells in a vascular endothelial growth factor (VEGF)-dependent manner (PubMed:23529610). .; [Isoform 1]: Cell membrane; Single-pass type I membrane protein. Cytoplasm. Nucleus. Cytoplasmic vesicle. Early endosome. Detected on caveolae-enriched lipid rafts at the cell surface. Is recycled from the plasma membrane to endosomes and back again. Phosphorylation triggered by VEGFA binding promotes internalization and subsequent degradation. VEGFA binding triggers internalization and translocation to the nucleus.; [Isoform 2]: Secreted .; [Isoform 3]: Secreted.

**Tissue specificity** Detected in cornea (at protein level). Widely expressed.


#### **Function**

Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., Function: Receptor for VEGF or VEGFC. Has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 7 Ig-like C2-type (immunoglobulin-like) domains., subunit: Interacts with MYOF (By similarity). Interacts with SHB; upon VEGF activation. Interacts with HIV-1 Tat...

### **I** Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using VEGFR2 (Phospho-Tyr1054) Antibody



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using VEGFR2 (Phospho-Tyr1054) Antibody. The picture on the right is blocked with the phospho peptide.

### | Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Flk-1/Flt-4 (Phospho Tyr1054/1063)
Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents