Applications

WB,IHC,IF,ELISA



MYPT1 (Phospho Thr696) Rabbit pAb

CatalogNo: YP1131

Key Features

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype
• 115kD (Calculated) IgG

Recommended Dilution Ratios

WB 1:500-2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:10000

Not yet tested in other applications.

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human MYPT1

around the phosphorylation site of Thr696. AA range:661-710

Specificity

Phospho-MYPT1 (T696) Polyclonal Antibody detects endogenous levels of MYPT1 protein only when phosphorylated at T696. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RStQG

Target Information

Gene name

PPP1R12A

Protein Name

Protein phosphatase 1 regulatory subunit 12A

Organism	Gene ID	UniProt ID
Human	<u>4659;</u>	<u>014974;</u>
Mouse	<u>17931;</u>	Q9DBR7;
Rat	<u>116670;</u>	Q10728;


Cellular Localization Cytoplasm . Cytoplasm, cytoskeleton, stress fiber . Also along actomyosin filaments. .

Tissue specificity Expressed in striated muscles, specifically in type 2a fibers (at protein level).


Function

Function: Regulates myosin phosphatase activity., PTM: Phosphorylated by CIT (Rhoassociated kinase) (By similarity). Phosphorylated cooperatively by ROCK1 and CDC42BP on Thr-696. Phosphorylated on upon DNA damage, probably by ATM or ATR., sequence Caution: Contaminating sequence. Potential poly-A sequence., similarity: Contains 6 ANK repeats., subcellular location: Along actomyosin filaments and stress fibers., subunit: PP1 comprises a catalytic subunit, PPP1CA, PPP1CB or PPP1CC, and one or several targeting or regulatory subunits. PPP1R12A mediates binding to myosin. Interacts with ARHA and CIT (By similarity). Binds PPP1R12B, ROCK1 and IL16.,

Validation Data



Immunofluorescence analysis of A549 cells, using MYPT1 (Phospho-Thr696) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using MYPT1 (Phospho-Thr696) Antibody. The picture on the right is blocked with the phospho peptide.

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
MYPT1 (Phospho
Thr696) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents