PIP5KIII (Phospho Ser307) Rabbit pAb

CatalogNo: YP1162

Key Features

Host Species
- Rabbit

Reactivity
- Human,Mouse,Rat

Applications
- IHC,IF,ELISA

MW
- 237kD (Calculated)

Isotype
- IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:100-1:300
IF 1:200-1:1000
ELISA 1:10000
Not yet tested in other applications.

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized peptide derived from human PIP5K around the phosphorylation site of Ser307. AA range:273-322
Specificity	Phospho-PIP5KIII (S307) Polyclonal Antibody detects endogenous levels of PIP5KIII protein only when phosphorylated at S307.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SAsIT

Target Information

Gene name PIKFYVE

Protein Name 1-phosphatidylinositol 3-phosphate 5-kinase


Organism	Gene ID	UniProt ID
Human	200576;	Q9Y2I7;
Mouse	18711;	Q9Z1T6;

Cellular Localization Endosome membrane ; Peripheral membrane protein . Early endosome membrane ; Peripheral membrane protein. Cytoplasmic vesicle , phagosome membrane ; Peripheral membrane protein . Late endosome membrane ; Peripheral membrane protein . Mainly associated with membranes of the late endocytic pathway. .


Tissue specificity Brain ,Epithelium ,PCR rescued clones ,T-cell ,

Function Catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4-phosphate = ADP + 1-phosphatidyl-1D-myo-inositol 4 ,5-bisphosphate. ,Disease:Defects in PIKFYVE are the cause of corneal fleck dystrophy (CFD) [MIM:121850]. CFD is an autosomal dominant disorder of the cornea characterized by numerous small white flecks scattered in all levels of the stroma. Although CFD may occasionally cause mild photophobia , patients are typically asymptomatic and have normal vision. ,Function:Supports the intracellular PIP pool and to a lesser extent , the PI 4 ,5-P (2) pool. It generates PIP from PI and , to a lesser extent , PI 4 ,5-P (2) from PI 4-P. There are indications that it phosphorylates the D-5 rather than the D-4 position. Has a role in endosome-related membrane trafficking. ,similarity:Contains 1 DEP domain. ,similarity:Contains 1 FYVE-type zinc finger. ,similarity:Contains 1 PI5K domain. ,subcellular location:Mainly associated with membranes of the late endocytic pathway. ,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PIP5K (Phospho-Ser307) Antibody



Immunofluorescence analysis of HeLa cells, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
PIP5KIII (Phospho Ser307) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.