

PIP5KIII (Phospho Ser307) Rabbit pAb

CatalogNo: YP1162

Key Features

Host Species

Rabbit

Reactivity

Human,Mouse,Rat

Applications

IHC,IF,ELISA

MW • 237kD (Calculated)

Isotype • IgG

Recommended Dilution Ratios

IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:10000 Not yet tested in other applications.

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information


ImmunogenThe antiserum was produced against synthesized peptide derived from human PIP5K
around the phosphorylation site of Ser307. AA range:273-322

Specificity Phospho-PIP5KIII (S307) Polyclonal Antibody detects endogenous levels of PIP5KIII protein only when phosphorylated at S307.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SAsIT


Target Information

Gene name	PIKFYVE			
Protein Name	1-phosphatidylinositol 3-phosphate 5-kinase			
	Organism	Gene ID	UniProt ID	
	Human	<u>200576;</u>	<u>Q9Y2I7;</u>	
	Mouse	<u>18711;</u>	<u>Q9Z1T6;</u>	
Cellular Localization	Endosome membrane ; Peripheral membrane protein . Early endosome membrane ; Peripheral membrane protein. Cytoplasmic vesicle, phagosome membrane ; Peripheral membrane protein . Late endosome membrane ; Peripheral membrane protein . Mainly associated with membranes of the late endocytic pathway			
Tissue specificity	Brain, Epithelium, PCR rescued clor	nes,T-cell,		
Function	of corneal fleck dystrophy (CFD) [the cornea characterized by nume stroma. Although CFD may occasi	bisphosphate.,Disease MIM:121850]. CFD is a erous small white fleck onally cause mild pho ision.,Function:Suppor It generates PIP from ions that it phosphory related membrane trat E-type zinc finger.,sim	e:Defects in PIKFYVE are the cause an autosomal dominant disorder of cs scattered in all levels of the tophobia, patients are typically ts the intracellular PIP pool and to a PI and, to a lesser extent, PI 4,5- lates the D-5 rather than the D-4 fficking.,similarity:Contains 1 DEP ilarity:Contains 1 PI5K	


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PIP5K (Phospho-Ser307) Antibody



Immunofluorescence analysis of HeLa cells, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using PIP5K (Phospho-Ser307) Antibody. The picture on the right is blocked with the phospho peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **PIP5KIII (Phospho Ser307) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents