**Applications** 

WB,IHC,IF,ELISA



# Integrin β2 (Phospho Thr758) Rabbit pAb

Reactivity

CatalogNo: YP1171 Orthogonal Validated 💽

## **Key Features**

Host Species

Rabbit
 Human, Mouse, Rat

MW Isotype
• 170kD (Observed) • IgG

## **Recommended Dilution Ratios**

WB 1:500-2000 IHC 1:50-300 IF 1:200-1:1000 ELISA 5000-1:20000

## **Storage**

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

# **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human

CD18/ITGB2 around the phosphorylation site of Thr758. AA range:720-769

#### **Specificity**

Phospho-Integrin β2 (T758) Polyclonal Antibody detects endogenous levels of Integrin β2 protein only when phosphorylated at T758. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):SAtTT

# **Target Information**

**Gene name** 

ITGB2

**Protein Name** 

Integrin beta-2

Organism	Gene ID	UniProt ID
Human	<u>3689;</u>	<u>P05107;</u>
Mouse	<u>16414;</u>	<u>P11835;</u>


Cellular Localization Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein.

**Tissue specificity** Leukocytes (PubMed:23775590). Expressed in neutrophils (at protein level) (PubMed:21193407, PubMed:28807980).


#### **Function**

Disease: Defects in ITGB2 are the cause of leukocyte adhesion deficiency type I (LAD1) [MIM:116920]. LAD1 patients have recurrent bacterial infections and their leukocytes are deficient in a wide range of adhesion-dependent functions., Function: Integrin alpha-L/beta-2 is a receptor for ICAM1, ICAM2, ICAM3 and ICAM4. Integrins alpha-M/beta-2 and alpha-X/beta-2 are receptors for the iC3b fragment of the third complement component and for fibrinogen. Integrin alpha-X/beta-2 recognizes the sequence G-P-R in fibrinogen alpha-chain. Integrin alpha-M/beta-2 recognizes P1 and P2 peptides of fibrinogen gamma chain. Integrin alpha-M/beta-2 is also a receptor for factor X. Integrin alpha-D/beta-2 is a receptor for ICAM3 and VCAM1., online information: ITGB2 mutation db, PTM: Both Ser-745 and Ser-756 become phosphorylated when T-cells are exposed to phorbol esters. Phosphorylation on Thr-758 (but not on Ser-756) allows interaction with 14-3-3 proteins., similarity: Belongs to the integrin beta chain family., similarity: Contains 1 VWFA domain., subunit: Heterodimer of an alpha and a beta subunit. Beta-2 associates with either alpha-L, alpha-M, alpha-X or alpha-D. Interacts with COPS5 and RANBP9.,


# **I** Validation Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using CD18/ITGB2 (Phospho-Thr758) Antibody



Immunofluorescence analysis of HepG2 cells, using CD18/ITGB2 (Phospho-Thr758) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from COS7 cells treated with EGF 200ng/ml 30', using CD18/ITGB2 (Phospho-Thr758) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

# | Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Integrin β2** 

(Phospho Thr758) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody   ELISA Kits   Protein   Reagents
	Immunoway - 4 / 4