

# Vimentin (Phospho Tyr38) Rabbit pAb

CatalogNo: YP1209 Orthogonal Validated 💽

## Key Features

Host Species <ul> <li>Rabbit</li> </ul>	Reactivity <ul> <li>Human:Y38,Mouse:Y38,Rat:Y38</li> </ul>	Applications • WB,ELISA
MW • 51kD (Observed)	Isotype • IgG	

#### **Recommended Dilution Ratios**

WB 1:500-10000 ELISA 1:10000

## **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	PBS, pH 7.4, containing 0.02% sodium azide as Preservative and 50% Glycerol.

#### **Basic Information**

Clonality Polyclonal

## Immunogen Information

ImmunogenSynthesized phospho-peptide around the phosphorylation site of human Vimentin<br/>(Phospho-Tyr38)SpecificityThis antibody detects endogenous levels of Vimentin at Human:Y38;Mouse:Y38;Rat:Y38,<br/>It doesn't reacte with total protein.The name of modified sites may be influenced by<br/>many factors, such as species (the modified site was not originally found in human<br/>samples) and the change of protein sequence (the previous protein sequence is<br/>incomplete, and the protein sequence may be prolonged with the development of protein<br/>sequencing technology). When naming, we will use the "numbers" in historical reference<br/>to keep the sites consistent with the reports. The antibody binds to the following<br/>modification sequence (lowercase letters are modification sites):RTySL

## Target Information

Gene	name	VIM	

#### Protein Name vimentin

Human	<u>7431;</u>	<u>P08670;</u>
Mouse	<u>22352;</u>	<u>P20152;</u>
Rat	<u>81818;</u>	<u>P31000;</u>

## Cellular

Cytoplasm . Cytoplasm, cytoskeleton . Nucleus matrix . Cell membrane .

- Localization
- **Tissue specificity** Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.
- Function Function: Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells., online information: Vimentin entry, PTM: One of the most prominent phosphoproteins in various cells of mesenchymal origin. Phosphorylation is enhanced during cell division, at which time vimentin filaments are significantly reorganized., sequence Caution: Intron retention., similarity: Belongs to the intermediate filament family., subunit: Homopolymer. Interacts with HCV core protein. Interacts with LGSN and SYNM., tissue specificity: Highly expressed in fibroblasts, some expression in T- and B-lymphocytes, and little or no expression in Burkitt's lymphoma cell lines. Expressed in many hormone-independent mammary carcinoma cell lines.,

## Validation Data



Western Blot analysis of HELA cells using Antibody diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Vimentin (Phospho-Ser38) Antibody



Western blot analysis of lysates from 293 cells treated with paclitaxel, using p-Vimentin (Phospho-Ser38) antibody.

## **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: Vimentin (Phospho Tyr38) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents