**Applications** 

WB



# ASK1 (Phospho Thr845) Rabbit pAb

CatalogNo: YP1265

# **| Key Features**

Host Species Reactivity

Rabbit
 Human,Rat,Mouse,

MW Isotype
• 155kD (Observed) • IgG

#### **Recommended Dilution Ratios**

WB 1:1000-2000

# Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

# **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

**Immunogen** Synthesized phosho peptide around human ASK1 (Thr845)

**Specificity** This antibody detects endogenous levels of Human ASK1 (phospho-Thr845)

## | Target Information

Gene name MAP3K5 ASK1 MAPKKK5 MEKK5

#### **Protein Name** ASK1 (Thr845)

| Or | ganism | Gene ID        | UniProt ID      |
|----|--------|----------------|-----------------|
| ŀ  | Human  | <u>4217;</u>   | Q99683;         |
| ı  | Mouse  | <u>26408</u> ; | <u>035099</u> ; |

Cellular Localization

Cytoplasm . Endoplasmic reticulum. Interaction with 14-3-3 proteins alters the distribution of MAP3K5/ASK1 and restricts it to the perinuclear endoplasmic reticulum region.

**Tissue specificity** Abundantly expressed in heart and pancreas.

**Function** 

Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Contains an N-terminal autoinhibitory domain. Activated by phosphorylation at Thr-838, inhibited by phosphorylation at Ser-966 and Ser-1033. Binds to, and stabilizes MAP3K6 and is activated by MAP3K6 by phosphorylation on Thr-838.,Function:Component of a protein kinase signal transduction cascade. Phosphorylates and activates MAP2K4 and MAP2K6, which in turn activate the JNK and p38 MAP kinases, respectively. Overexpression induces apoptotic cell death.,induction:By TNF-alpha. Inhibited by HIV-1 Nef.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Homodimer when inactive. Binds both upstream activators and downstream substrates in multimolecular complexes. Associates with and inhibited by HIV-1 Nef. Interacts with DAB2IP and PPM1L.,tissue specificity:Abundantly expressed in heart and pancreas.,

#### Validation Data

## Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **ASK1 (Phospho** 

Thr845) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents