

SIRT1 (Phospho Ser27) Rabbit pAb

CatalogNo: YP1495

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA, IHC

MW

- 85-110kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

IHC 1:50-300

ELISA 1:2000-20000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho peptide around human SIRT1 (Ser27)

Specificity This antibody detects endogenous levels of Human SIRT1 (phospho-Ser27)

Target Information

Gene name SIRT1 SIR2L1

Protein Name

SirT1 (Ser27)

Organism	Gene ID	UniProt ID
Human	23411 ;	Q96EB6 ;
Mouse	93759 ;	Q923E4 ;


Cellular Localization

Nucleus, PML body . Cytoplasm . Nucleus . Recruited to the nuclear bodies via its interaction with PML (PubMed:12006491). Colocalized with APEX1 in the nucleus (PubMed:19934257). May be found in nucleolus, nuclear euchromatin, heterochromatin and inner membrane (PubMed:15469825). Shuttles between nucleus and cytoplasm (By similarity). Colocalizes in the nucleus with XBP1 isoform 2 (PubMed:20955178). .; [SirT1 75 kDa fragment]: Cytoplasm . Mitochondrion .

Tissue specificity Widely expressed.

Function

Catalytic activity:NAD(+) + an acetylprotein = nicotinamide + O-acetyl-ADP-ribose + a protein.,cofactor:Binds 1 zinc ion per subunit.,enzyme regulation:Inhibited by nicotinamide. Activated by resveratrol (3,5,4'-trihydroxy-trans-stilbene), butein (3,4,2',4'-tetrahydrochalcone), piceatannol (3,5,3',4'-tetrahydroxy-trans-stilbene), Isoliquiritigenin (4,2',4'-trihydroxychalcone), fisetin (3,7,3',4'-tetrahydroxyflavone) and quercetin (3,5,7,3',4'-pentahydroxyflavone). RPS19BP1/AROS acts as a positive regulator of deacetylation activity.,Function:NAD-dependent deacetylase, which regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. Deacetylates 'Lys-382' of p53/TP53 and impairs its ability to induce proapoptotic program and modulate cell senescence. Deacetylates TAF1B and thereby represses rDNA transcription by the RNA polymerase I. Involved in HES1- and HEY2-mediated transcriptional repression. Inhibits skeletal muscle differentiation by deacetylating PCAF and MYOD1. May serve as a sensor of the cytosolic ratio of NAD(+)/NADH, which is essential in skeletal muscle cell differentiation. Despite some ability to deacetylate histones in vitro, such activity is either weak or inexistent in vivo. In case of HIV-1 infection, interacts with and deacetylates the viral Tat protein.,miscellaneous:Red wine, which contains resveratrol, may participate in activation of sirtuin proteins, and may therefore participate in an extended lifespan as it has been observed in yeast.,similarity:Belongs to the sirtuin family.,similarity:Contains 1 deacetylase sirtuin-type domain.,subcellular location:Recruited to the nuclear bodies via its interaction with PML.,subunit:Interacts with TAF1B. Found in a complex with PCAF and MYOD1 (By similarity). Interacts with MLLT7/FOXO4, HES1, HEY2, p53/TP53 and PML. Interacts with RPS19BP1/AROS.,tissue specificity:Widely expressed.,

Validation Data

Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
SIRT1 (Phospho Ser27) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)