

eNOS (Phospho Ser114) Rabbit pAb

CatalogNo: YP1844 Orthogonal Validated 💽

Key Features

Host Species

Rabbit

Reactivity

Human, Mouse, Rat

Applications

IHC,WB

MW

• 130kD (Observed)

Recommended Dilution Ratios

WB 1:500-2000 IHC 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human eNOS (Phospho Ser114)

Specificity

This antibody detects endogenous levels of eNOS (Phospho Ser114) Rabbit pAb at Human, Mouse,Rat.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RPsPG

Target Information

Gene name

NOS3

Protein Name

Nitric oxide synthase, endothelial (Constitutive NOS) (cNOS) (EC-NOS) (Endothelial NOS) (eNOS) (NOS type III) (NOSIII)

Organism	Gene ID	UniProt ID
Human	<u>4846</u> ;	<u>P29474;</u>
Mouse	<u>18127;</u>	<u>P70313;</u>
Rat	24600;	<u>Q62600;</u>

Cellular Localization


Cell membrane. Membrane, caveola. Cytoplasm, cytoskeleton. Golgi apparatus. Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity.

Tissue specificity Platelets, placenta, liver and kidney.

Function

Catalytic activity:L-arginine + n NADPH + n H(+) + m H(+) + m H(+) = citrulline + nitric oxide + n NADP(+).,cofactor:Binds 1 FAD.,cofactor:Binds 1 FMN.,cofactor:Heme group.,cofactor:Tetrahydrobiopterin (BH4). May stabilize the dimeric form of the enzyme., enzyme regulation: Stimulated by calcium/calmodulin. Inhibited by NOSIP and NOSTRIN...Function:Produces nitric oxide (NO) which is implicated in vascular smooth muscle relaxation through a cGMP-mediated signal transduction pathway. NO mediates vascular endothelial growth factor (VEGF)-induced angiogenesis in coronary vessels and promotes blood clotting through the activation of platelets., online information: Nitric oxide synthase entry, polymorphism: Variation in NOS3 seem to be associated with susceptibility to coronary spasm., similarity: Belongs to the NOS family., similarity: Contains 1 FAD-binding FRtype domain., similarity: Contains 1 flavodoxin-like domain., subcellular location: Specifically associates with actin cytoskeleton in the G2 phase of the cell cycle; which is favored by interaction with NOSIP and results in a reduced enzymatic activity., subunit: Homodimer. Interacts with NOSIP and NOSTRIN., tissue specificity: Platelets, placenta, liver and kidney.,

I Validation Data



Western Blot analysis of 1 HEK-293 cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

eNOS (Phospho Ser114) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents