

www.immunoway.com

Bim (Phospho Thr56/116) Rabbit pAb

CatalogNo: YP1860 Orthogonal Validated 💽

Key Features

Host Species

Rabbit

Reactivity

Human,Mouse,Rat

Applications
• IHC,WB

MW • 22kD (Calculated)

Recommended Dilution Ratios

WB 1:500-2000 IHC 1:50-200

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human Bim (Phospho Thr56/116)

Specificity This antibody detects endogenous Bim(EL) only when phosphorylated at thr116 and endogenous Bim(L) when phosphorylated at thr56.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):TQtPS

Target Information

Gene name	BCL2L11 BIM			
Protein Name	Bcl-2-like protein 11 (Bcl2-L-11) (Bcl2-interacting mediator of cell death)			
	Organism	Gene ID	UniProt ID	
	Human	<u>10018;</u>	<u>043521;</u>	
	Mouse	<u>12125;</u>	<u>054918;</u>	
	Rat	<u>64547;</u>	<u>088498;</u>	
Cellular Localization	Endomembrane system ; Peripheral membrane protein . Associated with intracytoplasmic membranes; [Isoform BimEL]: Mitochondrion. Translocates from microtubules to mitochondria on loss of cell adherence.; [Isoform BimL]: Mitochondrion.; [Isoform BimS]: Mitochondrion.; [Isoform Bim-alpha1]: Mitochondrion.			
Tissue specificity	Isoform BimEL, isoform BimL and isoform BimS are the predominant isoforms and are widely expressed with tissue-specific variation. Isoform Bim-gamma is most abundantly expressed in small intestine and colon, and in lower levels in spleen, prostate, testis, heart, liver and kidney.			
Function	Bim-alpha2 and isoform Bim- isoforms BimEL, BimL and Bir to the Bcl-2 family.,subcellula	pre potent than isoform Biml alpha3 induce apoptosis, alt nS. Isoform Bim-gamma ind r location:Associated with ir	EL. Isoform Bim-alpha1, isoform hough less potent than the uces apoptosis.,similarity:Belongs	

membranes.,subunit:Forms heterodimers with a number of antiapoptotic Bcl-2 proteins including MCL1, BCL2, BCL2L1 isoform Bcl-X(L), BCL2A1/BFL-1, and BHRF1. Does not heterodimerize with proapoptotic proteins such as BAD, BOK, BAX or BAK.,tissue specificity:Isoform BimEL, isoform BimL and isoform BimS are the predominant isoforms and are ubiquitously expressed with a tissue-specific variation. Isoform Bim-gamma is most abundantly expressed in small intestine and colon, and in lower levels in spleen, prostate, testis, heart, liver and kidney.,

Validation Data



Western Blot analysis of 1 Jurkat cell, 2 Serum-free treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Bim (Phospho Thr56/116) Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents