**Applications** 

IHC,IF,ELISA



# **AKAP 250 Rabbit pAb**

CatalogNo: YT0166

### **Key Features**

Host Species Reactivity

Rabbit
 Human,Rat,Mouse,

MW Isotype
• 191kD (Calculated) • IgG

#### Recommended Dilution Ratios

IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:20000

Not yet tested in other applications.

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human AKAP12.

AA range:301-350

**Specificity** AKAP 250 Polyclonal Antibody detects endogenous levels of AKAP 250 protein.

## **Target Information**

Gene name AKAP12

**Protein Name** 

A-kinase anchor protein 12

Organism	Gene ID	UniProt ID	
Human	<u>9590;</u>	<u>Q02952;</u>	
Mouse		Q9WTQ5;	


Cellular Localization Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Membrane; Lipid-anchor. May be part of the cortical cytoskeleton.

**Tissue specificity** Expressed in endothelial cells, cultured fibroblasts and osteosarcoma, but not in platelets, leukocytes, monocytic cell lines or peripherical blood cells.


**Function** 

Caution: The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data., Disease: Antibodies to the C-terminal of gravin can be produced by patients with myasthenia gravis (MG).,Domain:Polybasic regions located between residues 266 and 557 are involved in binding PKC., Function: Anchoring protein that mediates the subcellular compartmentation of protein kinase A (PKA) and protein kinase C (PKC), induction: Activated by lysophosphatidylcholine (lysoPC)..PTM:Phosphorvlated upon DNA damage, probably by ATM or ATR., similarity: Contains 3 AKAP domains., subcellular location: May be part of the cortical cytoskeleton., subunit: Binds to dimeric RII-alpha regulatory subunit of PKC., tissue specificity: Expressed in endothelial cells, cultured fibroblasts and osteosarcoma, but not in platelets, leukocytes, monocytic cell lines or peripherical blood cells.,

### Validation Data



Immunofluorescence analysis of HeLa cells, using AKAP12 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using AKAP12 Antibody. The picture on the right is blocked with the synthesized peptide.

### Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **AKAP 250 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents