**Applications** 

WB,IHC,IF,ELISA



# **AKAP 79 Rabbit pAb**

CatalogNo: YT0167

### **Key Features**

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype • 47kD (Observed) IgG

### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:10000

Not yet tested in other applications.

#### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

# **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human AKAP5.

AA range:1-50

**Specificity** AKAP 79 Polyclonal Antibody detects endogenous levels of AKAP 79 protein.

## **Target Information**

Gene name

AKAP5

**Protein Name** 

A-kinase anchor protein 5

Organism	Gene ID	UniProt ID	
Human	<u>9495</u> ;	<u>P24588;</u>	
Mouse		D3YVF0;	


Cellular Localization Postsynaptic recycling endosome membrane; Lipid-anchor. Associates with lipid rafts...

Tissue specificity Predominantly in the cerebral cortex and the postsynaptic densities of the forebrain, and to a lesser extent in adrenal medulla, lung and anterior pituitary.


**Function** 

Domain:RII-alpha binding site, predicted to form an amphipathic helix, could participate in protein-protein interactions with a complementary surface on the R-subunit dimer., Function: May anchor the PKA protein to cytoskeletal and/or organelle-associated proteins, targeting the signal carried by cAMP to specific intracellular effectors. Association with to the beta2-adrenergic receptor (beta2-AR) not only regulates beta2-AR signaling pathway, but also the activation by PKA by switching off the beta2-AR signaling cascade., miscellaneous: The N-terminal region, which is highly basic, is required for interaction with calmodulin., similarity: Contains 1 AKAP domain., subcellular location: Associated with particulate fractions., subunit: Binding protein for dimer of the RIIbeta regulatory subunit of cAMP-dependent protein kinase (PKA) and also for the protein kinase C (PKC) and the phosphatase calcineurin (PP2B). Each enzyme is inhibited when bound to the anchoring protein. Also binds the beta2-adrenergic receptor., tissue specificity: Predominantly in the cerebral cortex and the postsynaptic densities of the forebrain, and to a lesser extent in adrenal medulla, lung and anterior pituitary.,


### Validation Data



Western Blot analysis of various cells using AKAP 79 Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of HeLa cells, using AKAP5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2, HeLa, and COLO205 cells, using AKAP5 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

### Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

AKAP 79 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody   ELISA Kits   Protein   Reagents
	Immunoway - 4 / 4