

www.immunoway.com

ARA70 Rabbit pAb

CatalogNo: YT0302 Orthogonal Validated 💽

Comparable Abs

Key Features

Host SpeciesRabbit

MW • 70kD (Observed) ReactivityHuman,Mouse,RatIsotype

• IgG

Applications

WB,IHC,IF,CoIP,ELISA

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000 IP 1:50-200Not yet tested in other applications IF 1:50-200

Storage

Storage*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

ImmunogenSynthesized peptide derived from ARA70 . at AA range: 320-400SpecificityARA70 Polyclonal Antibody detects endogenous levels of ARA70 protein.

Target Information

Gene name	NCOA4		
Protein Name	Nuclear receptor coactivator 4 Organism	Gene ID	UniProt ID
	Human	<u>8031;</u>	<u>Q13772;</u>
Cellular Localization	nucleus,autolysosome,		
Tissue specificity	Widely expressed. Also detected in adipose tissues and in different cell lines. Isoform Beta is only expressed in testis.		
Function	Alternative products:Additional isoforms seem to exist,Disease:A chromosomal aberration involving NCOA4 is a cause of thyroid papillary carcinoma (PACT) [MIM:188550]. Inversion inv(10)(q11.2;q11.2) generates the RET/NCOA4 (PTC3) oncogene that has been found in sporadic and radiation-associated post-Chernobyl thyroid papillary carcinomas.,Function:Enhances the androgen receptor transcriptional activity in prostate cancer cells. Ligand-independent coactivator of the peroxisome proliferator-activated receptor (PPAR) gamma.,polymorphism:Polymorphisms are specific to isoform alpha. Not found in isoform beta. Could be encoded by two different genes.,subunit:Interacts with the androgen receptor and the retinoid X receptor (RXR) in a ligand-dependent manner.,tissue specificity:Widely expressed. Also detected in adipose tissues and in different cell lines. Isoform beta is only expressed in testis.,		

Validation Data



Western Blot analysis of various cells using ARA70 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

	293	
		250
		150
		100
		75
NCOA4		
		50
		37
		25
		20
		20
		15
		(kd)
		(

Western blot analysis of NCOA4 Antibody. The lane on the right is blocked with the NCOA4 peptide.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **ARA70 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents