

c-FLIP Rabbit pAb

CatalogNo: YT0877

| Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, IHC, IF, ELISA

MW

- 55kD (Observed)

Isotype

- IgG

| Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

ELISA 1:20000

IF 1:50-200

| Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality

Polyclonal

| Immunogen Information

Immunogen

The antiserum was produced against synthesized peptide derived from human internal CFLAR. AA range: 181-230


Specificity

c-FLIP Polyclonal Antibody detects endogenous levels of c-FLIP protein.

| Target Information

Gene name	CFLAR		
Protein Name	CASP8 and FADD-like apoptosis regulator		
	Organism	Gene ID	UniProt ID
	Human	8837 ;	O15519 ;
	Mouse		O35732 ;
Cellular Localization	cytoplasm,cytosol,death-inducing signaling complex,CD95 death-inducing signaling complex,membrane raft,riposome,		
Tissue specificity	Widely expressed. Higher expression in skeletal muscle, pancreas, heart, kidney, placenta, and peripheral blood leukocytes. Also detected in diverse cell lines. Isoform 8 is predominantly expressed in testis and skeletal muscle.		
Function	<p>Domain:The caspase domain lacks the active sites residues involved in catalysis.,Function:Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6 mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase) activity.,induction:Repressed by IL-2 after TCR stimulation, during progression to the S-phase of the cell cycle.,PTM:Proteolytically processed; probably by caspase-8. Processing likely occurs at the DISC and generates subunit p43 and p12.,similarity:Belongs to the peptidase C14A family.,similarity:Contains 2 DED (death effector) domains.,subunit:TNFRSF6 stimulation triggers recruitment to the death-inducing signaling complex (DISC) formed by TNFRSF6, FADD and caspase-8. A proteolytic fragment (p43) stays associated with the DISC. Also interacts with caspase-10, caspase-3, TRAF1, TRAF2 and Bcl-X(L) (in vitro). Interacts with HBV protein X.,tissue specificity:Widely expressed. Higher expression in skeletal muscle, pancreas, heart, kidney, placenta, and peripheral blood leukocytes. Also detected in diverse cell lines. Isoform 8 is predominantly expressed in testis and skeletal muscle.,</p>		

Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
c-FLIP Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)