

Hexb Rabbit pAb

CatalogNo: YT2128

Key Features

Host Species

- Rabbit

Reactivity

- Human,Rat,Mouse,

Applications

- WB,IHC,IF,ELISA

MW

- 63kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-1:2000

IHC 1:100-1:300

IF 1:200-1:1000

ELISA 1:20000

Not yet tested in other applications.

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

The antiserum was produced against synthesized peptide derived from human HEXB. AA range:481-530

Specificity

Hexb Polyclonal Antibody detects endogenous levels of Hexb protein.

| Target Information

Gene name HEXB

Protein Name Beta-hexosaminidase subunit beta


Organism	Gene ID	UniProt ID
Human	3074 ;	P07686 ;
Mouse		P20060 ;

Cellular Localization Lysosome . Cytoplasmic vesicle, secretory vesicle, Cortical granule .


Tissue specificity Liver,Skin,

Function Catalytic activity:Hydrolysis of terminal non-reducing N-acetyl-D-hexosamine residues in N-acetyl-beta-D-hexosaminides.,Disease:Defects in HEXB are the cause of GM2-gangliosidosis type 2 (GM2G2) [MIM:268800]; also known as Sandhoff disease. GM2-gangliosidosis is an autosomal recessive lysosomal storage disease marked by the accumulation of GM2 gangliosides in the neuronal cells. GM2G2 is clinically indistinguishable from GM2-gangliosidosis type 1, presenting startle reactions, early blindness, progressive motor and mental deterioration, macrocephaly and cherry-red spots on the macula.,Function:Responsible for the degradation of GM2 gangliosides, and a variety of other molecules containing terminal N-acetyl hexosamines, in the brain and other tissues.,online information:HEXB mutation database,PTM:N-linked glycans at Asn-142 and Asn-190 consist of Man(3)-GlcNAc(2) and Man(5 to 7)-GlcNAc(2), respectively.,PTM:The beta-A and beta-B chains are produced by proteolytic processing of the precursor beta chain.,similarity:Belongs to the glycosyl hydrolase 20 family.,subunit:There are 3 forms of beta-hexosaminidase: hexosaminidase A is a trimer composed of one subunit alpha, one subunit beta chain A and one subunit beta chain B; hexosaminidase B is a tetramer of two subunit beta chains A and two subunit beta chains B; hexosaminidase S is an homodimer of two alpha subunits. The two beta chains are derived from the cleavage of the beta subunit.,

| Validation Data



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using HEXB Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using HEXB Antibody. The lane on the right is blocked with the synthesized peptide.

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Hexb Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)