

Trk B Rabbit pAb

CatalogNo: YT4743

| Key Features

Host Species

Rabbit

MW

• 145kd or 92kd(with limited samples) (Observed)

Reactivity

Human, Mouse, Rat

Isotype
• IgG

Applications

• WB,IHC,IF,ELISA

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:20000 IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Trk B. AA

range:671-720

Specificity Trk B Polyclonal Antibody detects endogenous levels of Trk B protein.

Target Information

Gene name

NTRK2

Protein Name

BDNF/NT-3 growth factors receptor

Organism	Gene ID	UniProt ID
Human	<u>4915;</u>	<u>Q16620;</u>
Mouse	<u>18212;</u>	<u>P15209;</u>
Rat	<u>25054;</u>	<u>Q63604;</u>

Cellular Localization

Cell membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein . Early endosome membrane . Cell projection, axon . Cell projection, dendrite. Cytoplasm, perinuclear region. Cell junction, synapse, postsynaptic density. Internalized to endosomes upon ligand-binding. .

Tissue specificity Isoform TrkB is expressed in the central and peripheral nervous system. In the central nervous system (CNS), expression is observed in the cerebral cortex, hippocampus, thalamus, choroid plexus, granular layer of the cerebellum, brain stem, and spinal cord. In the peripheral nervous system, it is expressed in many cranial ganglia, the ophthalmic nerve, the vestibular system, multiple facial structures, the submaxillary glands, and dorsal root ganglia. Isoform TrkB-T1 is mainly expressed in the brain but also detected in other tissues including pancreas, kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in the brain.

Function

Alternative products:Additional isoforms seem to exist, Catalytic activity: ATP + a [protein]-Ltyrosine = ADP + a [protein]-L-tyrosine phosphate., Function: Receptor for brain-derived neurotrophic factor (BDNF), neurotrophin-3 and neurotrophin-4/5 but not nerve growth factor (NGF). Involved in the development and/or maintenance of the nervous system. This is a tyrosine-protein kinase receptor. Known substrates for the TRK receptors are SHC1, PI-3 kinase, and PLC-gamma-1.,PTM:Ligand-mediated auto-phosphorylation.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 2 Ig-like C2-type (immunoglobulin-like) domains., similarity: Contains 2 LRR (leucine-rich) repeats..subunit:Exists in a dynamic equilibrium between monomeric (low affinity) and dimeric (high affinity) structures. Binds SH2B2. Interacts with SQSTM1 and KIDINS220., tissue specificity: Isoform TrkB is widely expressed, mainly in the nervous tissue. In the CNS, expression is observed in the cerebral cortex, hippocampus, thalamus, choroid plexus, granular layer of the cerebellum, brain stem, and spinal cord. In the peripheral nervous system, it is expressed in many cranial ganglia, the ophtalmic nerve, the vestibular system, multiple facial structures, the submaxillary glands, and dorsal root ganglia. Isoform TrkB-T1 is expressed in multiple tissues, mainly in brain, pancreas, kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in brain.,

I Validation Data

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Trk B Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents