

ADCYA Rabbit pAb

CatalogNo: YT6551

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 177kD (Calculated)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-2000

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human ADCYA AA range: 368-418

Specificity This antibody detects endogenous levels of ADCYA at Human/Mouse/Rat

Target Information

Gene name ADCY10 SAC

Protein Name ADCYA

| Organism | Gene ID | UniProt ID |
|----------|--------------------------|--------------------------|
| Human | 55811 ; | Q96PN6 ; |
| Mouse | 271639 ; | Q8C0T9 ; |
| Rat | 59320 ; | Q9Z286 ; |

Cellular Localization

Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm, cytoskeleton . Cytoplasm, perinuclear region . Nucleus . Cell projection, cilium . Cytoplasm . Mitochondrion . Distributed to subcellular compartments containing cAMP targets. Found as a plasma membrane-associated protein, protein concentrated in the perinuclear region and protein colocalized with actin or tubulin. .


Tissue specificity

Detected in airway epithelial cells and testis (at protein level) (PubMed:17591988). Weakly expressed in multiple tissues. Expressed in brain, heart, kidney, liver, lung, pancreas, peripheral blood leukocytes, placenta, skeletal muscle, stomach, thymus, airway epithelial cells, duodenum, jejunum and ileum. Very low level of expression in bone.

Function

Catalytic activity:ATP = 3',5'-cyclic AMP + diphosphate.,cofactor: Binds 2 magnesium ions per subunit.,Disease:Genetic variations in ADCY10 are associated with absorptive hypercalciuria type 2 (HCA2) [MIM:143870]. Absorptive hypercalciuria (AH) is a common cause of calcium oxalate nephrolithiasis. Clinically, AH is characterized by intestinal hyperabsorption of calcium in the presence of normal serum calcium and immunoreactive PTH (iPTH). It is often accompanied by low bone mineral density (BMD), particularly of the lumbar spine. About 50% of patients with AH present with a family history of calcium oxalate nephrolithiasis and hypercalciuria.,enzyme regulation:Activated by manganese or magnesium ions. In the presence of magnesium ions, the enzyme is activated by bicarbonate while in the presence of manganese ions, the enzyme is inhibited by bicarbonate. In the absence of magnesium and bicarbonate, the enzyme is weakly activated by calcium.,Function:Soluble adenylyl cyclase that has a critical role in mammalian spermatogenesis. Produces the cAMP which mediates in part the cAMP-responsive nuclear factors indispensable for maturation of sperm in the epididymis. Induces capacitation, the maturational process that sperm undergo prior to fertilization. May be the bicarbonate sensor. Involved in ciliary beat regulation.,PTM: Cleavage may occur to generate the active 48 kDa form.,similarity:Belongs to the adenylyl cyclase class-4/guanylyl cyclase family.,similarity:Contains 2 guanylate cyclase domains.,subcellular location:Distributed to subcellular compartments containing cAMP targets. Found as a plasma membrane-associated protein, protein concentrated in the perinuclear region and protein colocalized with actin or tubulin.,tissue specificity:Weakly expressed in multiple tissues. Expressed in brain, heart, kidney, liver, lung, pancreas, peripheral blood leukocytes, placenta, skeletal muscle, stomach, thymus, airway epithelial cells, duodenum, jejunum and ileum. Very low level of expression in bone.,

| Validation Data



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°C over night

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
ADCYA Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)