

# **NOSTN Rabbit pAb**

CatalogNo: YT7453

# Key Features

Host Species

Rabbit

Reactivity

Human,Mouse,Rat

Applications
• WB

MW • 56kD (Calculated) Isotype • IgG

# **Recommended Dilution Ratios**

#### WB 1:500-2000

### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

Clonality Polyclonal

### Immunogen Information

Immunogen	Synthesized peptide derived from human NOSTN AA range: 309-359
Specificity	This antibody detects endogenous levels of NOSTN at Human/Mouse/Rat


### **Target Information**

Gene name NOSTRIN

#### Protein Name NOSTN

	Organism	Gene ID	UniProt ID	
	Human	<u>115677;</u>	<u>Q8IVI9;</u>	
	Mouse	<u>329416;</u>	<u>Q6WKZ7;</u>	
	Rat	<u>311111;</u>	<u>Q5I0D6;</u>	
Cellular Localization	[Isoform 1]: Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle . Cytoplasm, cytoskeleton . Enriched in selected actin structures (PubMed:16234328, PubMed:16376344).; [Isoform 3]: Nucleus .			
Tissue specificity	Expressed at highest levels in heart, kidney, placenta and lung, and at lowest levels in brain, thymus and spleen. Present in vascular endothelial cells and placenta. Over-expressed in placenta from women with pre-eclampsia (at protein level).			
Function	Domain:The FCH domain is necessary for membrane targeting.,Domain:The SH3 domain mediates interaction with NOS3, DNM2 and WASL.,Function:Multivalent adapter protein which may decrease NOS3 activity by inducing its translocation away from the plasma membrane.,similarity:Contains 1 FCH domain.,similarity:Contains 1 SH3 domain.,subcellular location:Enriched in selected actin structures.,subunit:Homotrimer. Interacts with DAB2 (By similarity). Interacts with NOS3, DNM2, WASL and CAV1.,tissue specificity:Expressed at highest levels in heart, kidney, placenta and lung, and at lowest levels in brain, thymus and spleen. Present in vascular endothelial cells and placenta. Over-expressed in placenta from women with pre-eclampsia (at protein level).,			

# Validation Data



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night

# **Contact information**

Orders:order@immunoway.comSupport:tech@immunoway.comTelephone:877-594-3616 (Toll Free), 408-747-0185Website:http://www.immunoway.comAddress:2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **NOSTN Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents