

# RN103 Rabbit pAb

CatalogNo: YT7532

### Key Features

Host Species

Rabbit

Reactivity

Human,Mouse,Rat

Applications
• WB,IHC

MW • 75kD (Calculated) Isotype • IgG

#### **Recommended Dilution Ratios**

WB 1:500-2000 IHC 1:50-300

#### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Polyclonal

#### Immunogen Information

Immunogen	Synthesized peptide derived from human RN103 AA range: 243-293
Specificity	This antibody detects endogenous levels of RN103 at Human/Mouse/Rat

#### Target Information

Gene name RNF103 ZFP103

#### Protein Name RN103

Organism	Gene ID	UniProt ID
Human	<u>7844;</u>	<u>000237;</u>
Mouse	<u>22644;</u>	<u>Q9R1W3;</u>
Rat		<u>Q9EPZ8;</u>

## Cellular

Endoplasmic reticulum membrane ; Multi-pass membrane protein .

#### Localization

**Tissue specificity** Highly expressed in the normal cerebellum but not in the cerebral cortex.

**Function** Domain: The acidic C-terminus and the basic N-termminus are thought to render the protein in a closed, soluble and inactive conformation through an autoinhibitory intramolecular interaction. The open and active conformation, which enables membrane binding and oligomerization, is achieved by interaction with other cellular binding partners, probably including other ESCRT components., Function: Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P2 and PtdIns(3,4)P2 in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Required for sorting/trafficking of EGF receptor., miscellaneous: Its overexpression strongly inhibits HIV-1 release., similarity: Belongs to the SNF7 family...similarity:Contains 1 RING-type zinc finger...subcellular location:Localizes to the midbody of dividing cells., subunit: Probable core component of the endosomal sorting required for transport complex III (ESCRT-III). ESCRT-III components are thought to multimerize to form a flat lattice on the perimeter membrane of the endosome. Several assembly forms of ESCRT-III may exist that interact and act sequentally. Forms a metastable monomer in solution; its core structure (without part of the putative autoinhibitory C-terminal acidic region) oligomerizes into a flat lattice via two different dimerization interfaces. In vitro, heteromerizes with CHMP2A (but not CHMP4) to form helical tubular structures that expose membrane-interacting sites on the outside whereas VPS4B can associate on the inside of the tubule. May interact with IGFBP7; the relevance of such interaction however remains unclear. Interacts with CHMP2A. Interacts with CHMP4A: the interaction requires the release of CHMP4A autoinhibition. Interacts with VPS4A. Interacts with STAMBP; the interaction appears to relieve the autoinhibition of CHMP3.,tissue specificity: Highly expressed in the normal cerebellum but not in the cerebral cortex.,tissue specificity:Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.,

#### Validation Data



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000,  $4^{\circ}$  over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

### **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **RN103 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents